

**PE Products collection  
of labels as  
Purchased by  
Plaintiffs**

**Sandra Yousefzadeh**

**- PE Products-**

**Sudafed PE® Sinus Pressure + Pain  
Maximum Strength**

**-Johnson & Johnson Consumer Inc.**



# **Anntwanette Jones**

## **-PE Products-**

**Equate Children's Multi-Symptom Cold  
Liquid, Very Berry  
-liquid**

**- Walmart, Inc.**



No Coating Area  
COATING FREE AREA

COATING FREE AREA

Glue

N.C.N

COATING FREE AREA  
No Coating Area

**Drug Facts**

**Active Ingredients (in each 5 mL) Purpose**  
Dextromethorphan HBr 5 mg.....Cough suppressant  
Guaifenesin 100 mg.....Expectorant  
Phenylephrine HCl 2.5 mg.....Nasal decongestant

**Uses**  
■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive  
■ temporarily relieves:  
■ cough due to minor throat and bronchial irritation as may occur with a cold or inhaled irritants  
■ the intensity of coughing  
■ the impulse to cough to help your child get to sleep  
■ nasal congestion due to a cold  
■ stuffy nose

**Warnings**  
Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has:  
■ heart disease  
■ high blood pressure  
■ thyroid disease  
■ diabetes  
■ persistent or chronic cough such as occurs with asthma  
■ cough that occurs with too much phlegm (mucus)

**When using this product**  
■ do not use more than directed

**Stop use and ask a doctor if**  
■ your child gets nervous, dizzy or sleepless  
■ symptoms do not get better within 7 days or occur with fever  
■ cough lasts more than 7 days, comes back, or occurs with fever, rash or persistent headache.  
These could be signs of a serious illness.

**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than 6 doses in any 24-hour period  
■ measure only with dosing cup provided  
■ do not use dosing cup with other products  
■ dose as follows or as directed by a doctor

**Drug Facts (continued)**

| Age                                | Dose                |
|------------------------------------|---------------------|
| children 6 years to under 12 years | 10 mL every 4 hours |
| children 4 years to under 6 years  | 5 mL every 4 hours  |
| children under 4 years             | do not use          |

■ mL = milliliter

**Other information**  
■ each 5 mL contains: potassium 5 mg, sodium 5 mg  
■ store between 15-30°C (59-86°F)  
■ do not refrigerate  
■ dosing cup provided

**Inactive ingredients** citric acid anhydrous, D&C red #33, dextrose, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, methylparaben, potassium sorbate, propyl gallate, propylene glycol, propylparaben, purified water, saccharin sodium, sodium hydroxide, sorbitol solution, sucralose, xanthan gum

**Questions? 1-888-287-1915**

**DISTRIBUTED BY: Wal-Mart Stores, Inc., Bentonville, AR 72716**  
\*This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Children's Mucinex® Multi-Symptom Cold.

**KEEP BOX FOR COMPLETE INFORMATION**  
DO NOT USE IF SEAL UNDER CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

 Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

 Recycle logo: PLASTIC BOTTLE, PLASTIC CUP

equate™

children's  
**MULTI-SYMPTOM  
COLD**

AGES 4 to 12 YEARS

Very Berry Flavored

NDC 49035-715-03

**Compare to Children's Mucinex® Multi-Symptom Cold active ingredients**

**children's MULTI-SYMPTOM COLD**

Dextromethorphan HBr 5 mg  
Cough Suppressant  
Guaifenesin 100 mg  
Expectorant  
Phenylephrine HCl 2.5 mg  
Nasal Decongestant

AGES 4 to 12 YEARS

☐ Relieves stuffy nose  
☐ Soothes cough  
☐ Relieves chest congestion  
☐ Thins and loosens mucus

■ Alcohol Free

**Very Berry Flavored**  
4 FL OZ (118 mL)

 Dosing Cup Included

equate™

children's  
**MULTI-SYMPTOM  
COLD**

AGES 4 to 12 YEARS

Very Berry Flavored

NDC 49035-715-03

**Compare to Children's Mucinex® Multi-Symptom Cold active ingredients**

**children's MULTI-SYMPTOM COLD**

Dextromethorphan HBr 5 mg  
Cough Suppressant  
Guaifenesin 100 mg  
Expectorant  
Phenylephrine HCl 2.5 mg  
Nasal Decongestant

AGES 4 to 12 YEARS

☐ Relieves stuffy nose  
☐ Soothes cough  
☐ Relieves chest congestion  
☐ Thins and loosens mucus

■ Alcohol Free

**Very Berry Flavored**  
4 FL OZ (118 mL)

 Dosing Cup Included

381137

6 81131 07834 4

EPD

NFR# 53041 REV 08/17

**Delsym® Children's Cough+ Cold Nighttime  
Berry Flavored**

**-liquid**

**-RB Health (US) LLC**



2450

**COUGH+  
COLD  
NIGHTTIME**

1.00

**Drug Facts****Active ingredients (in each 10 mL)**

**Acetaminophen 325 mg...Pain reliever/fever reducer**  
Diphenhydramine HCl 12.5 mg...Antihistamine/  
cough suppressant  
Phenylephrine HCl 5 mg...Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - runny nose
  - sneezing
- temporarily reduces fever
- controls cough to help your child get to sleep

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on the skin
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if the child is**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**Drug Facts (continued)****When using this product**

- do not use more than directed (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or last more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL in dosing cup provided every 4 hours while symptoms last; do not give more than 5 doses in any 24-hour period
- children under 6 years of age: do not use

**Other information**

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)
- do not refrigerate

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate\*, xanthan gum \*may contain this ingredient

**Questions? 1-866-682-4639**

You may also report side effects to this phone number.

**Keep carton for full information.**

**Night**  
NDC 63824-212-64

**children's  
Delsym®**

AGES  
6+

**COUGH+  
COLD  
NIGHT+TIME**

Acetaminophen (Pain Reliever/Fever Reducer)  
Diphenhydramine HCl (Antihistamine/Cough Suppressant)  
Phenylephrine HCl (Nasal Decongestant)

- Relieves Cough, Stuffy Nose & Sore Throat
- Reduces Fever

Please visit our website  
[www.delsym.com](http://www.delsym.com)

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



Dosing cup included



**berry**  
Flavored Liquid

**4 fl oz (118 mL)**

Dist. by: RB Health (US), Parsippany, NJ 07054-0224  
©2022 RB Health 062322 3234388

1.00

**Vicks® Dayquil™ SEVERE Honey Cold &  
Flu**

**-Liquid Medicine**

**-Procter & Gamble Company**



| Drug Facts (continued)  |                   |                                 |                   |                            |                   |                           |              |                      |            |
|---|-------------------|---------------------------------|-------------------|----------------------------|-------------------|---------------------------|--------------|----------------------|------------|
| <ul style="list-style-type: none"><li>• pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)</li><li>• fever gets worse or lasts more than 3 days</li><li>• redness or swelling is present</li><li>• new symptoms occur</li><li>• cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.</li></ul> <p><b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> |                   |                                 |                   |                            |                   |                           |              |                      |            |
| Directions  |                   |                                 |                   |                            |                   |                           |              |                      |            |
| <ul style="list-style-type: none"><li>• take only as directed</li><li>• only use the dose cup provided</li><li>• do not exceed 4 doses per 24 hrs</li></ul> <table><tr><td>adults &amp; children 12 yrs &amp; over</td><td>30 mL every 4 hrs</td></tr><tr><td>children 6 to under 12 yrs</td><td>15 mL every 4 hrs</td></tr><tr><td>children 4 to under 6 yrs</td><td>ask a doctor</td></tr><tr><td>children under 4 yrs</td><td>do not use</td></tr></table>   |                   | adults & children 12 yrs & over | 30 mL every 4 hrs | children 6 to under 12 yrs | 15 mL every 4 hrs | children 4 to under 6 yrs | ask a doctor | children under 4 yrs | do not use |
| adults & children 12 yrs & over   | 30 mL every 4 hrs |                                 |                   |                            |                   |                           |              |                      |            |
| children 6 to under 12 yrs  | 15 mL every 4 hrs |                                 |                   |                            |                   |                           |              |                      |            |
| children 4 to under 6 yrs   | ask a doctor      |                                 |                   |                            |                   |                           |              |                      |            |
| children under 4 yrs  | do not use        |                                 |                   |                            |                   |                           |              |                      |            |
| Other information   |                   |                                 |                   |                            |                   |                           |              |                      |            |
| <ul style="list-style-type: none"><li>• each 15 mL contains: sodium 14 mg</li><li>• store at no greater than 25°C and do not refrigerate</li></ul>  |                   |                                 |                   |                            |                   |                           |              |                      |            |
| Inactive ingredients  |                   |                                 |                   |                            |                   |                           |              |                      |            |
| citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum  |                   |                                 |                   |                            |                   |                           |              |                      |            |
| Questions?  |                   |                                 |                   |                            |                   |                           |              |                      |            |
| 1-800-362-1683  |                   |                                 |                   |                            |                   |                           |              |                      |            |

| Drug Facts (continued)  |  |
|---|--|
| Warnings  |  |
| <b>Liver warnings:</b> This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"><li>• adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product</li><li>• child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product</li><li>• taken with other drugs containing acetaminophen</li><li>• adult has 3 or more alcoholic drinks every day while using this product</li></ul> <b>Allergy Alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • hives • rash If a skin reaction occurs, stop use and seek medical help right away.<br><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. |  |
| Do not use  |  |
| <ul style="list-style-type: none"><li>• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li><li>• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li></ul>  |  |
| Ask a doctor before use if you have   |  |
| <ul style="list-style-type: none"><li>• liver disease</li><li>• heart disease</li><li>• thyroid disease</li><li>• diabetes</li><li>• trouble urinating due to enlarged prostate gland</li><li>• cough that occurs with too much phlegm (mucus)</li><li>• persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li></ul>  |  |
| Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.   |  |
| When using this product, do not use more than directed.   |  |
| Stop use and ask a doctor if  |  |
| <ul style="list-style-type: none"><li>• you get nervous, dizzy or sleepless</li></ul>   |  |



# **Daniel Calzado**

## **-PE Products-**

**CVS Health Non-Drowsy Nasal Decongestant  
PE Maximum Strength  
-tablets**

**-CVS Pharmacy Inc.**







**CVS Health Non-Drowsy Sinus PE Pressure,  
Pain + Cold**

**-CVS Pharmacy Inc.**

**Drug Facts (continued)**

**Other information** ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

■ see end flap for expiration date and lot number

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Drug Facts (continued)**

**Inactive ingredients** corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, povidone, silicon dioxide, stearic acid

**Questions or comments?** 1-800-426-9391

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 325 mg  
Pain reliever/fever reducer

Phenylephrine HCl 5 mg  
Nasal decongestant

**Purpose**

Acetaminophen 325 mg  
Pain reliever/fever reducer

Phenylephrine HCl 5 mg  
Nasal decongestant

**Uses**

■ temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:  
■ sore throat and pain ■ nasal congestion ■ headache

■ promotes sinus drainage ■ temporarily reduces fever

**Warnings**

■ Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

**Directions**

■ do not use more than directed

■ adults and children 12 years and over

■ take 2 caplets every 4 hours

■ do not take more than 10 caplets in 24 hours

■ children under 12 years: ask a doctor

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Stop use and ask a doctor if**

■ fever gets worse or lasts more than 3 days

■ pain or nasal congestion gets worse or lasts more than 7 days

■ nervousness, dizziness, or sleeplessness occur

■ redness or swelling is present ■ new symptoms occur

These could be signs of a serious condition.

**Ask a doctor before use if you have**

■ liver disease ■ diabetes ■ thyroid disease ■ heart disease ■ difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product do not exceed recommended dosage.**

**Drug Facts (continued)**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not use more than directed

■ adults and children 12 years and over

■ take 2 caplets every 4 hours

■ do not take more than 10 caplets in 24 hours

■ children under 12 years: ask a doctor

#343398



Lot/Exp area at minimum size  
1.25 x .5"

CVS Health.

Non-Drowsy

Sinus PE Pressure + Pain

CVS Health.

Non-Drowsy

Sinus PE Pressure + Pain

ACETAMINOPHEN - Pain reliever, Fever reducer

PHENYLEPHRINE HYDROCHLORIDE - Nasal decongestant

MAXIMUM STRENGTH

Relieves:

- Sinus pain & headache
- Sinus pressure
- Nasal & sinus congestion

24 CAPLETS UNCOATED



Actual Size

CVS Quality  
Money Back Guarantee

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-0231-502-08-H  
REV0820050208

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark SUDAFED PE® SINUS PRESSURE + PAIN.  
REB0820050208  
Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2021 CVS/pharmacy CVS.com®  
1-800-SHOP CVS  
V-19849

**Vicks NyQuil™ SEVERE Maximum  
Strength Cold & Flu Berry Flavored  
-Liquid Medicine**

**-Procter & Gamble Company**



**Advil Multi-Symptom Cold & Flu  
-tablets**

**-Haleon U.S. Holdings L.L.C.**  
**(f/k/a GSK Consumer Healthcare Holdings (US) LLC)**





The image shows the packaging for Advil Multi-Symptom Cold & Flu. The top half is yellow with the brand name 'Advil' in large red letters. Below this is a blue band with the text 'MULTI-SYMPTOM COLD & FLU' in white. The bottom half is yellow and lists the ingredients: Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID), Phenylephrine HCl 10 mg Nasal Decongestant, and Chlorpheniramine Maleate 4 mg Antihistamine. To the right of the ingredients is a list of symptoms treated, each preceded by a red checkmark. A red banner in the top right corner says '50% MORE 10 FREE Tablets'. On the right side, a blue banner says 'ONE TABLET DOSE!'. At the bottom right, a white coated tablet is shown with 'Advil CF' printed on it, and the text '30/20 Coated Tablets' is displayed in large red and white numbers.

# Advil

**50% MORE  
10 FREE  
Tablets**

## MULTI-SYMPTOM COLD & FLU

**Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)  
Phenylephrine HCl 10 mg Nasal Decongestant  
Chlorpheniramine Maleate 4 mg Antihistamine**

- ✓ Headache
- ✓ Fever
- ✓ Body Aches
- ✓ Sinus Pressure
- ✓ Nasal Congestion
- ✓ Runny Nose
- ✓ Sneezing
- ✓ Nasal Swelling

**ONE  
TABLET  
DOSE!**

**30/20 Coated  
Tablets**

Advil  
CF



**Eli Erlick**

**-PE Products-**

**Acetaminophen Day/Night Time Vapor Ice  
Cold and Flu Relief Caplets - 24ct - up &  
up™**

**-tablets**


**-Target Brands Inc.**



**TYLENOL® Cold + Flu Severe For Day And Night Time**

**-**  
**—**

**-Johnson & Johnson Consumer**



123665

NDC 50580-414-04

**TYLENOL®** COLD + FLU  
FOR ADULTS  
**SEVERE**

**COLD + FLU  
SEVERE**

**DO NOT USE IF BLISTER UNIT IS TORN OR BROKEN**

NDC 50580-414-04

# **TYLENOL<sup>®</sup>**

**FOR ADULTS**

## **COLD + FLU**

## **SEVERE**

## Acetaminophen.

**Dextromethorphan HBr,  
Phenylephrine HCl, Guaifenesin**  
Pain Reliever-Fever Reducer, Cough Suppressant,  
Nasal Decongestant, Expectorant

## Acetaminophen

**Chlorpheniramine Maleate,  
Dextromethorphan HBr, Phenylephrine HCl  
Pain Reliever—Fever Reducer, Antihistamine,  
Cough Suppressant, Nasal Decongestant**

 DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION

Actual  
Size

## 16 CAPLETS

**★ NIGHT**

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



Actual  
Size

## 8 CAPLETS

**TOTAL 24 CAPLETS**

## PARENTS:

**Learn about teen medicine abuse**

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

30047050  
123665

How can we help?  
1-877-895-3665



## Drug Facts (continued)

**Sore throat warning:** If sore throat is severe or followed by fever, headache, rash, nausea

**Do not use**

- with any other drug containing acetaminophen are not sure whether a drug contains acetaminophen
- if you are now taking a prescription medication for depression, psychiatric or emotional
- 2 weeks after stopping the MAOI drug, it contains an MAOI, ask a doctor or pharmacist
- if you have ever had an allergic reaction

**Ask a doctor before use if you have**

- liver disease
- diabetes
- trouble urinating due to prostate problems
- persistent or chronic cough such as occurs with smoking
- cough that occurs with too much phlegm
- a breathing problem such as emphysema

**When using this product**

- do not exceed recommended dose
- excitability may occur, especially if chills
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may
- avoid alcoholic drinks
- be careful when driving a motor vehicle

### Drug Facts (continued)

- Stop use and ask a doctor if:
- new onsets, dizziness, or sleeplessness
  - pain, nasal congestion or cough gets worse
  - ever gets worse or lasts more than 3 days
  - new symptoms occur
  - cough or mucus
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health care provider about use. Keep out of reach of children.**

Center right away. (1-800-222-1222) Quick m  
as for children event if you do not notice any si

### Directions

- |  |  |
|--|--|
| adults and children<br>12 years and over | <ul style="list-style-type: none"> <li>■ take 2 capsules</li> <li>■ swallow</li> </ul> |
|--|--|

### Other information

- do not use if blister unit is torn or broken

**Inactive ingredients:** anhydrous citric acid, flavor, butyramide, mannitol, sodium citrate, sodium phosphate, sodium saccharin, sucralose, xanthan gum.

- polyethylene glycol, polysorbate 80, potassium sodium benzoate, sodium citrate dihydrate sodium

**Questions or comments? call 1-877-**



ea, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

phen (prescription or nonprescription). If you have phen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain antidepressants, or Parkinson's disease), or if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

to this product or any of its ingredients

**High blood pressure** ■ thyroid disease  
■ enlarged prostate gland  
■ asthma or emphysema  
■ chronic cough  
■ glaucoma  
■ diabetes  
■ heart disease  
■ trouble urinating due to an enlarged prostate gland  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
■ cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache at last

**FLU SEVERE NIGHT**

**TYLENOL® COLD + FLU SEVERE DAY**

### Drug Facts (continued)

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-272-1222) Quick medical attention is critical for adults as well as children under 12 years.

ask a doctor

ask a doctor

ask a doctor

ask a doctor

ask a doctor

Tylenol® Cold + Flu Severe Day Made in Italy  
Distributed by: JOHNSON & JOHNSON CONSUMER INC.  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA

©J&JCI 2020



**Important: Read all product information before using. Keep this box for important information.**  
Do not take the day and night caplets at the same time. Do not take more than a total of 10 caplets in a 24-hour period. Take only as directed.

### TYLENOL® COLD + FLU SEVERE NIGHT

### TYLENOL® COLD + FLU SEVERE DAY

**Drug Facts**

| Active ingredients (in each caplet) | Purpose                     |
|-------------------------------------|-----------------------------|
| Acetaminophen 325 mg                | Pain reliever/fever reducer |
| Chlorpheniramine maleate 2 mg       | Antihistamine               |
| Dextromethorphan HBr 10 mg          | Cough suppressant           |
| Phenylephrine HCl 5 mg              | Nasal decongestant          |

**Uses** ■ for the temporary relief of the following cold/flu symptoms:

- minor aches and pains
- headache
- sore throat
- nasal congestion
- cough
- sinus congestion and pressure
- sneezing and runny nose
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Drug Facts**

| Active ingredients (in each caplet) | Purpose                     |
|-------------------------------------|-----------------------------|
| Acetaminophen 325 mg                | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg          | Cough suppressant           |
| Guaifenesin 200 mg                  | Expectorant                 |
| Phenylephrine HCl 5 mg              | Nasal decongestant          |

**Uses** ■ for the temporary relief of the following cold/flu symptoms:

- minor aches and pains
- headache
- sore throat
- nasal congestion
- cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

LIFT HERE



**Vicks NyQuil™ Severe Cold & Flu,  
Maximum Strength**

**-Liquicaps**

**-Procter & Gamble Company**



**Drug Facts (continued)**

**Directions**

- take only as directed
- adults & children 12 yrs & over: 2 LiquiCaps with water every 4 hrs
- children 4 to under 12 yrs: ask a doctor
- children under 4 yrs: do not use

**Other information**

- store at no greater than 25°C

**Drug Facts (continued)**

**Inactive ingredients** FD&C Blue No. 1, gelatin, glycerin, pharmaceutical ink\*, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, sorbitan solution

\*may contain this ingredient

**Questions?** 1-800-362-1683

**MAX STRENGTH**

**VICKS®**

# NyQuil™

## SEVERE

### COLD & FLU

Acetaminophen, Phenylephrine HCl,  
Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat,  
Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

**24 LIQUICAPS™**

Nighttime Relief

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**TAMPER EVIDENT:**  
This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

**Made in Canada**  
DST. BY  
PROCTER & GAMBLE  
CINCINNATI OH 45202

**P&G**  
[www.pg.com](http://www.pg.com)  
[www.vicks.com](http://www.vicks.com)  
Patents:  
[www.pg.com/patents](http://www.pg.com/patents)  
90760628

**COATED PAPER, MULTI-LAYER BOX TRAY**  
This is required in all containers

**Drug Facts**

| Active ingredients (in each LiquiCap) | Purpose                     |
|---------------------------------------|-----------------------------|
| Acetaminophen 325 mg                  | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg            | Cough suppressant           |
| Doxylamine succinate 6.25 mg          | Antihistamine               |
| Phenylephrine HCl 5 mg                | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores free breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product**

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Vicks DayQuil™ and NyQuil™ VapoCOOL  
SEVERE Maximum Strength Cold & Flu**

**-Caplets**

**-Procter & Gamble Company**





## Daytime Vaporice™ Cold &amp; Flu

## ents (in each caplet)

| Active ingredients (in each caplet) | Purpose                   |
|-------------------------------------|---------------------------|
| Acetaminophen 325 mg                | Pain relief/fever reducer |
| Dextromethorphan HBr 10 mg          | Cough suppressant         |
| Quafenesin 200 mg                   | Expectorant               |
| Phenylephrine HCl 5 mg              | Nasal decongestant        |

**Uses** temporarily relieves common cold/flu symptoms: ■ fever ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ headache ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breath through the nose ■ minor aches and pains ■ helps

ocus) and thin bronchial secretion to rid the

- promotes nasal and sinus drainage ▼

1012

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks everyday while using this product.  
Allergy alert: Acetaminophen may cause severe skin reactions.

**Symptoms may include:** skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. Stop this at once if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, or consult a doctor promptly.

**Do not use** with any other drug containing acetaminophen (see cautions on prescription). If you are not sure whether adrug

contains acetaminophen, ask a doctor or pharmacist. **■** If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotion disorders, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **■** If you are prescribed an alternate medication.

Ask a doctor before use if you have [www.thirdspace.com](http://www.thirdspace.com) or diabetes

Ask, a do you therefore use if you have: ☐ dry food disease ☐ diabetes  
☐ liver disease ☐ heart disease ☐ high blood pressure ☐ trouble  
urinating due to an enlarged prostate gland ☐ cough that occurs with  
too much phlegm (mucus) ☐ persistent or chronic cough such as  
occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not use more than directed.

Stop use and ask a doctor if you have serious dizziness or lightheadedness.

■ pain nasal congestion, or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache if it lasts. This could be signs of a serious condition. ▼

**NTS:**  
medicine abuse  
lineAbuse.org

**PARENTS**  
Learn about test  
[www.StopMed.com](http://www.StopMed.com)

Fluor yellow 26 aluminum blue, fuchs malachite, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, sodium disulfate, stearic acid, sucralose, talc, titanium dioxide

\_\_\_\_\_

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-2122). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

■ do not exceed 8 caplets per 24 hrs

|                                 |                                  |
|---------------------------------|----------------------------------|
| adults & children 12 yrs & over | 2 caplets with water every 4 hrs |
| children 4 to under 12 yrs      | ask a doctor                     |
| children under 4 yrs            | do not use                       |

■ store at 20°-25°C (68°-77°F)

1





**John Slughter**

**-PE Products-**

**Vicks DayQuil™ SEVERE Honey Flavored  
Maximum Strength Cold & Flu  
-liquid**

**-Procter & Gamble Company**



**Drug Facts (continued)**

- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

**Questions?** 1-800-362-1683

**Drug Facts (continued)**

**Warnings**

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • blisters • rash If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless

Tamper Evident:  
Do not use if printed shrinkband seal  
around the neck is broken or missing.  
← **PEEL BACK FOR DRUG FACTS**

**Drug Facts**

| Active ingredients (in each 15 mL) | Purpose                     |
|------------------------------------|-----------------------------|
| Acetaminophen 325 mg               | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg         | Cough suppressant           |
| Guaifenesin 200 mg                 | Expectorant                 |
| Phenylephrine HCl 5 mg             | Nasal decongestant          |

**Uses**

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache • fever • sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

**Parents:**  
Learn about teen medicine risks  
www.StopMedicineAbuse.org

**P&G**  
www.pg.com

**PLASTIC BOTTLE**

91708228

12390004162

**Keith Mortuiccio**

**-PE Products-**



**TYLENOL® Cold + Flu Severe**

**-caplets**

**-Johnson & Johnson Consumer Inc.**



### Drug Facts

Active ingredients

Purpose

### Drug Facts (continued)

■ temporarily reduces fever



### Active ingredients (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/  
fever reducer  
Dextromethorphan HBr 20 mg.....Cough suppressant  
Guaifenesin 400 mg.....Expectorant  
Phenylephrine HCl 10 mg.....Nasal decongestant

### Uses

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - nasal congestion
  - sore throat
  - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

### Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

### Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

### Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dose**

**Stop use and ask a doctor if**

### Drug Facts (continued)

- redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

away. (1-800-222-1222)  
Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

#### Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

|                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | <ul style="list-style-type: none"> <li>■ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li> <li>■ do not take more than 150 mL in 24 hours, unless directed by a doctor</li> </ul> |
| children under 12 years               | ask a doctor   |

### Drug Facts (continued)

#### Other information

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

#### Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

### Drug Facts (continued)

#### Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**TYLENOL®**  
COLD + FLU SEVERE

How can we help?  
**1-877-895-3665**

Acetaminophen  
(650 mg per 30 mL)  
Dextromethorphan HBr  
(20 mg per 30 mL)  
Guaifenesin  
(400 mg per 30 mL)  
Phenylephrine HCl  
(10 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
© J&JCI 2021

30049845



**Mucinex Maximum Strength Sinus-Max®  
Day & Night**

**-liquid gels**

**-RB Health (USA) LLC**





**Drug Facts****Active ingredients (in each liquid gel) Purposes**  
**Mucinex SINUS-MAX DAY**

Acetaminophen 325 mg.....Pain reliever  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Guaifenesin 200 mg.....Expectorant  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Active ingredients (in each liquid gel) Purposes**  
**Mucinex SINUS-MAX NIGHT**

Acetaminophen 325 mg.....Pain reliever  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Doxylamine succinate 6.25 mg.....Antihistamine  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - headache
  - cough
  - minor aches and pains
  - sinus congestion and pressure
  - runny nose and sneezing (**NIGHT only**)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**DAY only**)

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- glaucoma (**NIGHT only**)

**Drug Facts (continued)**

■ a breathing problem such as emphysema or chronic bronchitis (**NIGHT only**)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**NIGHT only**)

**When using this product** ■ do not use more than directed

■ excitability may occur, especially in children (**NIGHT only**)

■ marked drowsiness may occur (**NIGHT only**)

■ alcohol, sedatives, and tranquilizers may increase drowsiness (**NIGHT only**)

■ avoid alcoholic drinks (**NIGHT only**)

■ be careful when driving a motor vehicle or operating machinery (**NIGHT only**)

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see **Overdose warning**)
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

**Other information**

- store at 20-25°C (68-77°F) ■ avoid excessive heat

**Inactive ingredients (DAY only)** FD&C yellow no. 6, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water

**Inactive ingredients (NIGHT only)** D&C yellow no. 10, FD&C blue no. 1, gelatin, glycerin, hypromellose, isopropyl alcohol, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, sorbitol sorbitan solution, titanium dioxide, water



**Mucinex**  
**SINUS-MAX**

MAXIMUM STRENGTH

Maximum Strength per 4-hour dose  
 Do not take the Mucinex Sinus-Max DAY and NIGHT liquid gels at the same time. Always wait at least 4 hours before taking another dose of Mucinex Liquid gel. Do not take more than a total of 12 liquid gels in a 24-hour period.  
 Take only as directed.  
 Keep carton for full information

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Mucinex Maximum Strength Fast-Max®  
Severe Congestion & Cough**



**-liquid**

**-RB Health (USA) LLC**





| MIN PT SIZE & HEIGHT (Non-Automated)    |  |       |  |
|---|--|-------|--|
| NET WT (6.3 FL OZ 180ml):               | .10inch  | 10pt  |  |
| Drug Facts Title:                       | .X0inch  | XXpt  |  |
| Active Ingredient Title:                | .07inch  | 7pt   |  |
| Active Ingredient (BOP Ingredients):    | .X0inch  | XXpt  |  |
| Min Size of all other Copy:             | .066inch   | 6.6pt |  |
| This Box<br>1/2 x 1/2<br>inch<br>Square | Front PDP in square inches :<br>12.136<br>Rear PDP in square inches :<br>XX.XX |       |  |

| RB Artwork and Print Specification   |  |                                 |   |                                  |  |               |               |               |               |
|--|--|---------------------------------|---|----------------------------------|--|---------------|---------------|---------------|---------------|
| Trident Reference No: RB267073<br>ZEN Ref: TR1237088<br>Action: A<br>Brand: Mucinex<br>Category: Adult<br>Segment Group: Fast Max<br>Segment: Severe Congestion Et Cough<br>Pack Size: 611 oz<br>Market/Country: USA<br>Date: 17/03/17   |  |                                 |   |                                  |  |               |               |               |               |
| RBH Contact: Laura Feld  |  |                                 |   |                                  |  |               |               |               |               |
| <b>Artwork Type: IDM Commercial</b><br>Component Code (as if applicable): 3048243<br>Parent Technical Packaging Specifications: D0319625<br>Finest Goods Code: 63924-01465-10<br>Supply Point: RB Hull<br>Pharmacode No/NE: N/A<br>Edgemark Position: NA   |  |                                 |   |                                  |  |               |               |               |               |
| CAD Cam Ref: Muc-Ub-D0319625-9047mm<br>Printer: Multi Packaging Solutions (Bourne, UK)<br>Substrate: Paper White   |  |                                 |   |                                  |  |               |               |               |               |
| <b>Technical &amp; Non Printing Items</b><br><table border="0"> <tr> <td><input type="checkbox"/> Cutter</td> <td><input type="checkbox"/> Cutter 2 (if applicable)</td> </tr> <tr> <td><input type="checkbox"/> Outline</td> <td><input type="checkbox"/> Outline 2 (if applicable)</td> </tr> </table> |  | <input type="checkbox"/> Cutter | <input type="checkbox"/> Cutter 2 (if applicable) | <input type="checkbox"/> Outline | <input type="checkbox"/> Outline 2 (if applicable) |               |               |               |               |
| <input type="checkbox"/> Cutter  | <input type="checkbox"/> Cutter 2 (if applicable)  |                                 |   |                                  |  |               |               |               |               |
| <input type="checkbox"/> Outline   | <input type="checkbox"/> Outline 2 (if applicable) |                                 |   |                                  |  |               |               |               |               |
| <b>Colours (Label Front)</b><br><table border="1"> <tr> <td>PAVING BLUE 1</td> <td>PAVING BLUE 2</td> <td>PAVING BLUE 3</td> <td>PAVING BLUE 4</td> </tr> <tr> <td>PAVING BLUE 5</td> <td>PAVING BLUE 6</td> <td>PAVING BLUE 7</td> <td>PAVING BLUE 8</td> </tr> </table>                                |  | PAVING BLUE 1                   | PAVING BLUE 2                                     | PAVING BLUE 3                    | PAVING BLUE 4                                      | PAVING BLUE 5 | PAVING BLUE 6 | PAVING BLUE 7 | PAVING BLUE 8 |
| PAVING BLUE 1  | PAVING BLUE 2                                      | PAVING BLUE 3                   | PAVING BLUE 4                                     |                                  |  |               |               |               |               |
| PAVING BLUE 5  | PAVING BLUE 6                                      | PAVING BLUE 7                   | PAVING BLUE 8                                     |                                  |  |               |               |               |               |
| <b>BARCODE INFO</b><br>Barcode Type: N/A<br>Barcode Number: N/A<br>Magnification: N/A<br>Truncated By: N/A<br>Full Height: N/A<br>Bar Height (Smallest Bar): N/A<br>BWR: N/A<br>Encoded Data: N/A  |  |                                 |   |                                  |  |               |               |               |               |
| <br>HEALTH > HYGIENE > HOME   |  |                                 |   |                                  |  |               |               |               |               |
| <br><b>TRIDENT</b><br>Connaught House, Connaught Road,<br>Kingswood Business Park, Hull, HU17 3AP, England.<br>T: +44 (0) 1482 828100   |  |                                 |   |                                  |  |               |               |               |               |
| Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.   |  |                                 |   |                                  |  |               |               |               |               |
| <b>STUDIO USE ONLY</b> Adam Simmons v2.0<br>Sm*Art check results: 0-1; 0-0; 0-0; - AS - 03/17/17 15:26:48  |  |                                 |   |                                  |  |               |               |               |               |
| <b>ARTWORK APPROVAL</b><br>Approved by Sonoco Trident on behalf of Recicltt Benckiser<br>Signed: _____<br>Date: _____  |  |                                 |   |                                  |  |               |               |               |               |
| <b>NOT APPROVED FOR COLOUR</b><br>UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.   |  |                                 |   |                                  |  |               |               |               |               |

**Vicks DayQuil™ and NyQuil™ VapoCOOL  
SEVERE Maximum Strength Cold & Flu +  
Congestion Relief Liquid Co-Pack**

**-Liquid**

**-Procter & Gamble Company**

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)

**CVS Health Daytime Cold & Flu Softgel**

**-tablets**

**-CVS Pharmacy Inc.**



2489



Compare to the active ingredients in  
Vicks® DayQuil® Severe Cold & Flu\*

**Severe**

**MAXIMUM STRENGTH**

**Daytime Softgels**

**Severe Cold/Flu Relief**

**ACETAMINOPHEN** - Pain reliever, Fever reducer  
**DEXTROMETHORPHAN HBr** - Cough suppressant  
**GUAIFENESIN** - Expectorant  
**PHENYLEPHRINE HCl** - Nasal decongestant

**Relieves:**  
Headache, Fever, Sore throat, Minor aches  
& pains, Cough, Chest congestion, Nasal/Sinus  
congestion & Sinus pressure

**Non-drowsy**  
**Alcohol free**  
**Antihistamine free**

  
Actual Size

**24 SOFTGELS**

\*This product is not  
manufactured or distributed  
by Procter & Gamble, owner  
of the registered trademarks  
Vicks® DayQuil® Severe Cold  
& Flu.

Distributed by:  
CVS Pharmacy, Inc.  
One CVS Drive  
Woonsocket, RI 02886  
© 2019 CVS/pharmacy  
CVS.com®  
1-800-SHOP CVS  
V-35248

  
CVS Quality  
Money Back Guarantee

**Drug Facts**

**Active ingredients (in each softgel)**

|                                 |                             |
|---------------------------------|-----------------------------|
| Acetaminophen 325 mg.....       | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg..... | Cough suppressant           |
| Guaifenesin 200 mg.....         | Expectorant                 |
| Phenylephrine HCl 5 mg.....     | Nasal decongestant          |

**Purpose**

**Drug Facts (continued)**

**Uses** ■ temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion & pressure ■ cough due to minor throat & bronchial irritation ■ minor aches & pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

**Warnings**  
**Liver warning** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product; ■ with other drugs containing acetaminophen; ■ 3 or more alcoholic drinks every day while using this product.  
**Allergy alert** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ take only as directed  
■ do not exceed 8 softgels per 24 hours

|                                   |                                     |
|-----------------------------------|-------------------------------------|
| adults & children 12 years & over | 2 softgels with water every 4 hours |
| children 4 to under 12 years      | ask a doctor                        |
| children under 4 years            | do not use                          |

**Other information** ■ store at room temperature

**Inactive ingredients** H14C Yellow # 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

**Questions or comments?**  
Call 1-888-333-9792

**THIS PRODUCT IS PACKAGED IN A CHILD-RESISTANT AND TAMPER-EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.**

**LOT NO.**

**EXP. DATE:**

#362042

50428161053



**Pedro Urena**

**- PE Products-**

**Alka-Seltzer Plus Severe Cold & Flu**

**-effervescent tablets**

**-Bayer HealthCare LLC**



**Theraflu Daytime Severe Cold Relief Berry  
Burst Flavor**

**-Hot Liquid Powder**

**-Haleon U.S. Holdings L.L.C.**  
**(f/k/a GSK Consumer Healthcare Holdings (US) LLC)**





**Vicks DayQuil™ SEVERE Maximum  
Strength Cold & Flu Daytime Relief  
-LiquiCaps**

**-Procter & Gamble Company**

**Drug Facts (continued)**

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs
- adults & children 12 yrs & over: 12 LiquiCaps with water every 4 hrs
- children 4 to under 12 yrs: ask a doctor
- children under 4 yrs: do not use

**Other information** store at no greater than 25°C

**Inactive Ingredients** FD&C Yellow No. 5, FD&C Yellow No. 6, gelatin, glycerin, lecithin, mica, polyethylene glycol, polyvinyl acetate phthalate, povidone, sorbitol sorbitan solution, titanium dioxide, water

**Questions? 1-800-362-1663**



**DayQuil**

**EASY TO OPEN BOTTLE<sup>1</sup>**

ACTUAL SIZE

**25% SMALLER\***

\*COMPARED TO ORIGINAL DAYQUIL™ 24Hr BUSTER PACK

<sup>1</sup>COMPARED TO ORIGINAL DAYQUIL™ 24Hr BUSTER PACK

**Drug Facts**

| Active ingredients (in each LiquiCap) | Purpose                     |
|---------------------------------------|-----------------------------|
| Acetaminophen 325 mg.....             | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg.....       | Cough suppressant           |
| Phenylephrine HCl 5 mg.....           | Nasal decongestant          |

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- hives
- itching

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland

TAMPER EVIDENT: DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED.

DIST. BY:  
**PROCTER & GAMBLE**  
CINCINNATI, OH 45202

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**P&G**  
[www.pg.com](http://www.pg.com)  
vicks.com  
Patients:  
[www.pg.com/patients](http://www.pg.com/patients)

Headache, Fever,  
Sore Throat,  
Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy

**25% SMALLER\***

**EASY TO SWALLOW**

**48 LIQUICAPS™**

**90491463**

**3 23900 03112 8**

**RECYCLED**

**PAPER BOX**

**PLASTIC BOTTLE**

©2013 P&G

**Kimberly McNulty**

**-PE Products-**

**up & up<sup>TM</sup> Daytime Severe Cold & Flu  
-Softgel**


**-Target Corporation**





**Walgreens Multi-Symptom Children's Cold  
-Liquid**

**- Walgreens**





Compare to Children's Mucinex® Multi-Symptom Cold active ingredients†

**children's**  
MULTI-SYMPTOM  
**Cold**

DEXTROMETHORPHAN HBr /  
COUGH SUPPRESSANT  
GUAIFENESIN / EXPECTORANT  
PHENYLEPHRINE HCl /  
NASAL DECONGESTANT


**ALCOHOL FREE**


- Relief from: stuffy nose, cough & chest congestion
- Breaks up mucus


**AGES**  
**4-12**  
**YEARS**



**4 FL OZ (118 mL)** **BERRY FLAVOR**







Compare to Children's Mucinex® Multi-Symptom Cold active ingredients†


**children's**  
MULTI-SYMPTOM  
**Cold**

DEXTROMETHORPHAN HBr /  
COUGH SUPPRESSANT  
GUAIFENESIN / EXPECTORANT  
PHENYLEPHRINE HCl /  
NASAL DECONGESTANT

**ALCOHOL FREE**


- Relief from: stuffy nose, cough & chest congestion
- Breaks up mucus

**AGES**  
**4-12**  
**YEARS**



**4 FL OZ (118 mL)** **BERRY FLAVOR**





**children's**  
MULTI-SYMPTOM  
**Cold**

DEXTROMETHORPHAN HBr /  
COUGH SUPPRESSANT  
GUAIFENESIN / EXPECTORANT  
PHENYLEPHRINE HCl /  
NASAL DECONGESTANT

**Drug Facts**

**Active Ingredients (in each 5 mL)**

Dextromethorphan HBr 5 mg.....Cough suppressant

Guaifenesin 100 mg.....Expectorant

Phenylephrine HCl 2.5 mg.....Nasal decongestant

**Purposes**

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

**Warnings**

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- your child gets nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- mL = milliliter

**Drug Facts (continued)**

| Age                                | Dose                |
|------------------------------------|---------------------|
| children 6 years to under 12 years | 10 mL every 4 hours |
| children 4 years to under 6 years  | 5 mL every 4 hours  |
| children under 4 years             | do not use          |

**Other Information**

- each 5 mL contains: sodium 3 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

**Inactive Ingredients** anhydrous citric acid, edetate disodium, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions or comments?**  
1-800-719-9260

**Gluten Free**

Walgreens Pharmacist Recommended  
Walgreens Pharmacist Survey  
†This product is not manufactured or distributed by Reckitt Benckiser, the distributor of Children's Mucinex® Multi-Symptom Cold.

**DOSAGE CUP PROVIDED**

**PARENTS:**  
Learn about teen medicine abuse.  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com ©2017 Walgreen Co.

ITEM 374549 W10106-1217-L



3 11917 17279 8

**LOT NO.**

**EXP.**

• 83926 94 C4

**Walgreens Daytime Severe Cold & Flu  
Maximum Strength**

**-Liquid**

**- Walgreens**



NDC 0363-5001-40

**Walgreens**Compare to the active ingredients in  
Vicks® DayQuil® Severe Cold & Flu™

DAYTIME • NON-DROWSY

# Severe Cold & Flu

**ACETAMINOPHEN / PAIN RELIEVER /  
FEVER REDUCER  
PHENYLEPHRINE HCl /  
NASAL DECONGESTANT  
DEXTROMETHORPHAN HBr /  
COUGH SUPPRESSANT  
GUAIFENESIN / EXPECTORANT****Maximum Strength**

- Headache, fever, sore throat, minor aches & pains
- Chest congestion, thins & loosens mucus
- Nasal/sinus congestion, sinus pressure
- Cough



Honey flavor

**12 FL OZ (355 mL)**

: 53U40 94 F1

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2022 Walgreen Co.**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)ITEM 714456  
W10107-0623-F  
REV-0323DO NOT USE IF PRINTED BACKGROUND  
IS BROKEN OR MISSING**Drug Facts****Active ingredients  
(in each 15 mL)**

| Active ingredients<br>(in each 15 mL) | Purpose                         |
|---------------------------------------|---------------------------------|
| Acetaminophen 325 mg.....             | Pain reliever/<br>fever reducer |
| Dextromethorphan HBr 10 mg.....       | Cough suppressant               |
| Guaifenesin 200 mg.....               | Expectorant                     |
| Phenylephrine HCl 5 mg.....           | Nasal decongestant              |

**Uses**

temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure  
■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat  
■ fever ■ cough due to minor throat and bronchial irritation  
■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**PEEL BACK AT  
CORNER FOR MORE  
INFORMATION**

: 53U40 94 B2

**Drug Facts  
(continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

HESIVE AREA  
NO VARNISH • NO TYPE**Drug Facts  
(continued)**

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

|                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |

ADHESIVE AREA  
TING • NO VARNISH • NO TYPE

2504

**AD**  
**NO COATING**

Ask a doctor before use if you have

- liver disease    ■ heart disease
- high blood pressure    ■ diabetes
- thyroid disease    ■ cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

**NO COATING**

|                      |            |
|----------------------|------------|
| children under 4 yrs | do not use |
|----------------------|------------|

**Other information** ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions? 1-800-719-9260**

\*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

\*\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® and DayQuil®.

**CVS Health Non-Drowsy Daytime Multi-Symptom Cold/Flu Relief**

**-Softgels**

**-CVS Pharmacy, Inc.**





**CVS Health Children's Day + Nighttime  
Cold, Cough, + Congestion Relief Liquid  
Combo Pack**

**-Liquid**

**-CVS Pharmacy, Inc.**



### Children's Cold & Cough Daytime

**Drug Facts**

**Active ingredients (in each 5 mL)**  
Dextromethorphan HBr 5 mg... Cough suppressant  
Phenylephrine HCl 2.5 mg... Nasal decongestant

**Uses**

- temporarily relieve
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

**Warnings**

**Do not use**

- in a child under 4 years of age
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- symptoms do not improve within 7 days or occur with fever
- cough persists more than 7 days, comes back or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not give more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- shake well before using

| Age                                 | Dose                |
|-------------------------------------|---------------------|
| children 6 to under 12 years of age | 10 mL every 4 hours |
| children 4 to under 6 years of age  | 5 mL every 4 hours  |
| children under 4 years of age       | do not use          |

### Children's Cold & Cough Daytime (continued)

**Drug Facts (continued)**

**Other information**

- each 5 mL contains sodium 2 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** acacia, dextrose, FD&C red #40, flavor, methyl, propylene glycol, purified water, sodium citrate

### Children's Nighttime Cold & Cough

**Drug Facts**

**Active ingredients (in each 5 mL)**  
Diphenhydramine HCl 12.5 mg... Antihistamine/Cough suppressant  
Phenylephrine HCl 2.5 mg... Nasal decongestant

**Uses**

- temporarily relieve
- runny nose
- itchy nose or throat
- itchy, watery eyes due to hay fever
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation as may occur with a cold

**Warnings**

**Do not use**

- in a child under 4 years of age
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- with any other product containing diphenhydramine, even one used or sold
- for the purpose of making your child sleepy

**Ask a doctor before use if the child has**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- chronic cough that lasts, or as occurs with asthma
- a breathing problem such as chronic bronchitis

**Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.**

### Children's Nighttime Cold & Cough (continued)

**Drug Facts (continued)**

**When using this product**

- do not exceed recommended dosage
- marked drowsiness may occur
- excitability may occur, especially in children
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- nausea, dizziness, or sleepiness occurs
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back, or occurs with a fever, rash, or persistent headache. These could be signs of a serious condition.

**Keep out of reach of children.**  
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not give more than 6 doses in 24 hours unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- shake well before using

| Age                                 | Dose                                   |
|-------------------------------------|--|
| children 6 to under 12 years of age | 10 mL every 4 hours                    |
| children 4 to under 6 years of age  | do not use unless directed by a doctor |
| children under 4 years of age       | do not use                             |

**Other information**

- each 5 mL contains sodium 3 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

**Inactive ingredients** acacia, dextrose, FD&C blue #1, FD&C red #40, flavor, methyl, propylene glycol, purified water, sodium benzoate, sodium citrate

**Questions or comments?**  
Call 1-877-763-3635 Monday-Friday 9AM-5PM EST

**PARENTS:**  
Learn about how to keep your child safe.  
[www.SuppliesHouse.org](http://www.SuppliesHouse.org)

## DAY & NIGHT COMBO PACK

**CVS Health.**

Compare to the active ingredients in Children's Triaminic® Day Time Cold & Cough\*

**Daytime**

### Children's Cold & Cough

DEXTROMETHORPHAN HBr  
Cough suppressant  
PHENYLEPHRINE HCl  
Nasal decongestant

**Relieves:**

- Cough
- Stuffy nose

**Alcohol 0.14%**

**For Ages 4 to 11 Years**

**Cherry Flavor**  
Dosage cup included  
Actual Bottle Size on Side Panel

**CVS Health.**

Compare to the active ingredients in Children's Triaminic® Night Time Cold & Cough\*

**Nighttime**

### Children's Nighttime Cold & Cough

DIPHENHYDRAMINE HCl  
Antihistamine; Cough suppressant  
PHENYLEPHRINE HCl  
Nasal decongestant

**Relieves:**

- Cough
- Runny, stuffy nose
- Itchy throat

**Alcohol free**

**For Ages 6 to 11 Years**

**Grape Flavor**  
Dosage cup included  
Actual Bottle Size on Side Panel

**4 FL OZ (118 mL) + 4 FL OZ (118 mL)  
TOTAL 8 FL OZ (236 mL)**

Package Contains Two Bottles  
Actual Size

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2019 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS V-13114



PLD-8319E  
PLD-8320E  
7/20/2019

Lot No.:  
Exp. Date:



#194300

**Tatyana Dekhtyar**

**-PE Products-**

**Advil Sinus Congestion & Pain**

**-tablets**

**-Haleon U.S. Holdings L.L.C.**

**(f/k/a GSK Consumer Healthcare Holdings (US) LLC)**

**NON-DROWSY**

# Advil

**50% MORE  
10 FREE Tablets**

## SINUS CONGESTION & PAIN

**Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)  
Phenylephrine HCl 10 mg Nasal Decongestant**

- ✓ Nasal Congestion
- ✓ Sinus Pressure
- ✓ Nasal Swelling
- ✓ Headache
- ✓ Fever
- ✓ Body Aches

**ONE TABLET DOSE!**

**30/20** Coated Tablets